

## The role of $11\beta$ -hydroxysteroid dehydrogenase in maturation of the intestine

I. Pohlová, I. Mikšík, J. Pácha \*

*Institute of Physiology, Czech Academy of Sciences, Vídeňská 1083,  
14220 Prague 4-Krč, Czech Republic*

Received 16 November 1996; received in revised form 22 April 1997

---

### Abstract

Glucocorticoids promote the development of many organs including intestine. At the cellular level, the activity of glucocorticoids is regulated by  $11\beta$ -hydroxysteroid dehydrogenase ( $11\beta$ HSD) which converts active glucocorticoids to inactive metabolites. As  $11\beta$ HSD is also expressed in the intestine, this enzyme may be an important regulator of intestinal maturation. To investigate this, we have performed the systematic study of the development of intestinal  $11\beta$ HSD activity and its cofactor preference as well as of the effect of  $11\beta$ HSD inhibition by carbenoxolone on postnatal development of sucrase, alkaline phosphatase and Na,K-ATPase in the intestine. The activity of  $11\beta$ HSD was low in ileum of suckling rats and significantly increased during the weaning period. In colon, the activity was already high in suckling rats and gradually rose during the postnatal development.  $11\beta$ HSD activity was undetectable in jejunum both in young and adult rats. At 14.5 nM corticosterone, colonic  $11\beta$ HSD utilized predominantly NAD as a cofactor, but displayed significant sensitivity also to NADP. Ileal  $11\beta$ HSD had similar sensitivity to both cofactors. With NAD as a cofactor, ileal  $11\beta$ HSD had a  $K_m$  ( $59 \pm 10$  nM) compatible with the colonic enzyme ( $81 \pm 14$  nM). Carbenoxolone administration to suckling and weanling rats *in vivo* did not result in any changes of sucrase activity in jejunum and ileum, alkaline phosphatase activity in ileum and distal colon or Na,K-ATPase activity in ileum. However, carbenoxolone significantly increased Na,K-ATPase activity in distal colon. Our results indicate that the high-affinity type of  $11\beta$ HSD is expressed not only in colon but also in ileum and that  $11\beta$ HSD is an important factor in the regulation of tissue levels of active glucocorticoids in developing colon but not in the small intestine. © 1997 Elsevier Science Ireland Ltd.

---

\* Corresponding author. Tel: +420 24752440; fax: +420 24719571.

**Keywords:** Mineralocorticoid receptors; Glucocorticoid receptors; Corticosterone; Steroid metabolism; Intestinal maturation;  $11\beta$ -Hydroxysteroid dehydrogenase; Carbenoxolone

---

## 1. Introduction

Postnatal maturation of the rodent intestine is characterized by significant modifications of enzyme activities and transport properties. These developmental changes in enzyme status and intestinal transport coincide with weaning period and are influenced by the hormonal milieu of the pups [1,2]. Several lines of evidence indicate that pituitary-adrenal system modulates intestinal development. Adrenalectomy or hypophysectomy of rats attenuate developmental changes and such alterations can be prevented by the administration of corticosterone [1]. Corticosterone, the plasma concentration of which increases just before the time of weaning [3] has been shown to affect the development of various enzymes including sucrase, alkaline phosphatase and peptidases in the small intestine [1,4]. Aldosterone, which has developmental profile similar to corticosterone, induces electrogenic amiloride-sensitive  $\text{Na}^+$  transport in suckling and weanling rats [5].

$11\beta$ -hydroxysteroid dehydrogenase ( $11\beta$ HSD), a microsomal enzyme responsible for the interconversion of glucocorticoids (corticosterone in the rat), to their inactive 11-dehydro metabolites (11-dehydrocorticosterone in the rat), is considered to be a potential regulator of glucocorticoid and mineralocorticoid action [6,7]. In glucocorticoid-target organs, it might modulate the access of endogenous glucocorticoids to glucocorticoid receptors via 11-dehydrogenase or 11-oxo-reductase pathway. In mineralocorticoid-target tissue, it protects the non-selective mineralocorticoid receptors from glucocorticoid excess. Two species of  $11\beta$ HSD exist: one is low-affinity and bidirectional ( $11\beta$ HSD1), the other is high-affinity and unidirectional ( $11\beta$ HSD2).  $11\beta$ HSD1 has higher  $K_m$  ( $\mu\text{M}$ ) for its glucocorticoid substrates, preference for NADP/NADPH as cofactors and seems to be an enzyme that produces active glucocorticoids from inactive metabolites in many tissues [8,9]. On the other hand,  $11\beta$ HSD2 is an enzyme, that has only  $11\beta$ -dehydrogenase activity, has nanomolar affinity for glucocorticoids and prefers NAD as cofactor [10]. The presence of both isoforms has been demonstrated in multiple organs including rat intestine [11,12]. The intestinal  $11\beta$ HSD activity is not distributed homogeneously along the intestine and its developmental profile is not identical in particular intestinal segments [13]. After birth  $11\beta$ HSD activity is high in caecum, proximal and distal colon and does not alter much until adulthood. Quite contrasting developmental profile is typical for ileum, where  $11\beta$ HSD activity is low during the first two postnatal weeks and increases during weaning period.

The aim of this study was to study cofactor dependence of  $11\beta$ HSD in the intestinal segments that express different developmental profiles of activity and to

investigate whether  $11\beta$ HSD plays a physiological role in the maturation of the intestinal epithelium. Using the  $11\beta$ HSD inhibitor, carbenoxolone [7], we examined the possible role of  $11\beta$ HSD in the developmental changes of sucrase, alkaline phosphatase and Na,K-ATPase activities during weaning. These enzymes were chosen because they are sensitive to corticosteroids and have distinct developmental profiles. Sucrase and Na,K-ATPase activities increase during development in the small and large intestine (1,14,15) whereas alkaline phosphatase decreases in ileum [16].

## 2. Materials and methods

### 2.1. Animals

Wistar rats were bred and maintained on a 12-h light: 12 h darkness cycle and received tap water ad libitum. The day of birth was designated day 0 and approximately 24 h after birth the litter size was reduced to 8 pups which were kept with the dams until they are 30 days old. To examine the effect of carbenoxolone on sucrase, alkaline phosphatase and Na,K-ATPase activities, each litter was divided into two groups. One group received i.p. injections of carbenoxolone (60 mg/kg body wt. day in 0.9% saline) or vehicle only (0.9% saline) from day 12 to day 23. Rats were killed by decapitation and segments of the intestine (jejunum, ileum, colon) were removed, rinsed with ice-cold saline, opened longitudinally and used for enzyme activity studies.

### 2.2. Assay of $11\beta$ HSD activity

The intestinal segments were homogenized in 0.2 M sucrose (1:4 w/v) using a Polytron homogenizer. The homogenate was centrifuged at  $1000 \times g$  for 10 min and the supernatant was assayed for protein concentration using the method of Bradford [17].  $11\beta$ HSD activity was determined by measuring the rate of conversion of corticosterone to 11-dehydrocorticosterone in assay tubes containing 250  $\mu$ l of intestinal homogenate (1 mg of protein), 750  $\mu$ l of buffer (100 mM KCl, 50 mM TRIS-HCl, pH 9.0) and 40  $\mu$ l NAD or NADP (final concentration 400  $\mu$ M). After 10 min preincubation at 37°C 1.3  $\mu$ Ci of [ $^3$ H]corticosterone (final concentration 14.5 nM) was added and the reaction continued 45 min. Preliminary studies indicated that using this amount of tissue protein the rate of reaction was linear within this time. The reaction was terminated by cooling of samples, which were then centrifuged 20 min ( $3000 \times g$ ). The supernatant was loaded onto C18 reverse phase Sep-Pak columns (Waters, Milford, MA, USA) and steroids were quantitatively (98%) eluted in 2 ml methanol. The samples were evaporated to dryness under nitrogen at 40°C, reconstituted in 100  $\mu$ l methanol, injected onto a steel cartridge (125  $\times$  4 mm internal diameter) packed with LiChrospher 100 RP-18 (5  $\mu$ m, Merck, Darmstadt, Germany) and analyzed by high performance liquid chromatographic system (Waters, Milford, MA, USA). The samples were separated

using a linear methanol-water gradient from 45:55 (v/v) to 65:35 (v/v) in 15 min and isocratic washing with 100% methanol for 10 min at a flow rate 1.0 ml/min. Column temperature was held at 45°C. The elution of  $^3\text{H}$ -labelled steroids were monitored by on-line radioactive detection using radioisotope detector with solid cell (Beckman Type 171, Fullerton, CA, USA). After subtraction of background radioactivity, integrated counts within peaks were analyzed by Apex Version 3.1 software (DataApex, Prague, Czech Republic). The activity of 11 $\beta$ HSD was expressed as the percentage distribution of radioactivity among the HPLC peaks of corticosterone and 11-dehydrocorticosterone in each chromatogram.

Dependence of 11 $\beta$ HSD activity on pH was measured as mentioned above using pH range of the buffer 7.0–9.5. Reaction was started after 10 min temperature equilibration by adding [ $^3\text{H}$ ]corticosterone and continued for another 45 min.

To determine the  $K_m$  of 11 $\beta$ HSD for corticosterone, aliquots of intestinal homogenates (ileum 250  $\mu\text{g}$  prot, colon 125  $\mu\text{g}$  prot) were incubated for 10 min (colon) or 20 min (ileum) at 37°C with 14.5 nM [ $^3\text{H}$ ]corticosterone and 0.0–1000 nM unlabeled corticosterone and fixed concentration of NAD (400  $\mu\text{M}$ ). The data were plotted according to the Lineweaver-Burk double-reciprocal linear transformation of the Michaelis-Menten equation and the least squares best fit computer program was used to calculate the  $K_m$  and  $V_{\text{max}}$  values.

### 2.3. Assay of sucrase

Sucrase was assayed in mid-jejunum and mid-ileum. For this purpose the segments were slit lengthwise and scraped with a steel spatula to remove mucosa. Homogenate of mucosa was prepared in 9 volumes of 154 mM KCl using a Teflon homogenizer. Sucrase activity was determined in 60 mM sodium maleate buffer (pH 6.0) with 300 mM sucrose at 37°C. The reaction was stopped by boiling and the glucose liberated was measured with the glucose oxidase reagent. Corrections were made for endogenous glucose in the tissue and in the substrate. Sucrase activity was expressed as  $\mu\text{moles}$  of glucose produced per hour and mg of protein. Protein was measured by the method of Bradford [17].

### 2.4. Assay of Na,K-ATPase

Na,K-ATPase activity was measured in crude homogenate as the ouabain-sensitive release of inorganic phosphate from ATP [14]. Briefly, the mucosa was scraped and homogenized with a Teflon pestle in nine volumes of the solution containing (mM): 30 TRIS-HCl; 250 sucrose; 5  $\text{Na}_2\text{EDTA}$ ; pH 7.3. Samples of homogenate were preincubated at 37°C for 10 min in a solution containing (in mM): 100 NaCl; 100 TRIS-HCl; 20 KCl, 5  $\text{MgCl}_2$ ; pH 7.3 with or without ouabain (final concentration 2 mM). The reaction was started by addition of ATP (final concentration 3.2 mM) and continued for 30 min. After stopping the reaction by trichloroacetic acid, the released inorganic phosphate was assayed. Na,K-ATPase activity was calculated as the difference between ATPase activity without and with ouabain and expressed as  $\mu\text{mol}$  of inorganic phosphate per mg of protein per hour.

### 2.5. Alkaline phosphatase

Alkaline phosphatase was measured according to Murer et al. [18]. Samples of homogenate were added to a mixture containing (in mM): 50 glycine-NaOH buffer (pH 10.5); 2 MgSO<sub>4</sub> and 2.5 ZnSO<sub>4</sub>. After a 10-min preincubation at 37°C the reaction was initiated by addition of p-nitrophenyl phosphate as substrate (final concentration 5.5 mM). The reaction was stopped with 2 M NaOH. The activity was expressed as  $\mu\text{mol}$  of released p-nitrophenol per mg of protein per hour.

### 2.6. Statistical analysis

The data are presented as the means  $\pm$  S.E.M. Group data were compared using Student's *t*-test. Statistical significance was set at  $P < 0.05$ .

## 3. Results

The developmental profile of 11 $\beta$ HSD activity (Fig. 1) and cofactor preference were investigated in homogenates of three intestinal segments - jejunum, ileum and colon. Colon is the site of a high 11 $\beta$ HSD activity which increases only moderately during development. In contrast, ileum of suckling rats has very low activity which is much higher in weaning period and adulthood. The conversion of corticosterone is absent in jejunum. At 14.5 nM corticosterone, the colonic 11 $\beta$ HSD has a clear preference for NAD over NADP activity in both young and adult animals. The enzyme used NADP to dehydrogenate corticosterone at 50% rate compared to that observed when NAD was used as a cofactor. The rate of conversion was very low in the absence of cofactors in all three segments. (Fig. 1). The effect of pH on the activity of 11 $\beta$ HSD is shown in Fig. 2. Unlike the NAD-dependent 11 $\beta$ HSD from cultured human colonic cells T84, which has a pH optimum of approximately seven [19], the rat colonic enzyme had a pH optimum of about nine. Carbenoxolone (0.1 M) strongly inhibited 11 $\beta$ HSD in both young ( $82 \pm 9\%$ ) and adult ( $88 \pm 10\%$ ) rats. The kinetic analysis of 11 $\beta$ HSD activity in ileum and colon revealed the presence of high-affinity 11 $\beta$ HSD in both intestinal segments (Table 1). Maximal capacity ( $V_{\text{max}}$ ) was, however, attenuated in ileum.

To investigate whether 11 $\beta$ HSD could play a role in the developmental changes of enzymes that are known to be sensitive to corticosteroids, we examined Na,K-ATPase and alkaline phosphatase activities in ileum and distal colon and sucrase activity in jejunum and ileum. As carbenoxolone was demonstrated to be efficient to inhibit 11 $\beta$ HSD (see above), the developmental changes of these enzymes were examined in the presence and absence of carbenoxolone in 24-day-old rats. The reason for the study of jejunum, which has no dehydrogenase activity (Fig. 1), was the finding of considerable 11-oxo-reductase activity in adult jejunum [20]. The data in Tables 2 and 3 show that carbenoxolone did not change neither the activity of alkaline phosphatase in colon and ileum nor the activity of sucrase in jejunum and ileum. In contrast to sucrase and alkaline phosphatase, carbenoxolone administra-




Fig. 1.  $11\beta$ -hydroxysteroid dehydrogenase activity in the presence of 14.5 nM corticosterone and its dependence on cofactors (NAD or NADP). Data are means  $\pm$  S.E.M. ( $n = 5$ ). The activity was expressed as conversion of corticosterone to 11-dehydrocorticosterone. Significantly different from the data obtained in the presence of NADP (\* $P < 0.05$ ).




Fig. 2. Effect of pH on activity of  $11\beta$ -hydroxysteroid dehydrogenase in distal colon of adult rats expressed as conversion of corticosterone to  $11$ -dehydrocorticosterone in the presence of NAD.

tion resulted in significant changes of distal colonic Na,K-ATPase but not of ileal Na,K-ATPase (Table 3). As shown in Fig. 3, the sucrase and Na,K-ATPase activities rose in intact rats during postnatal development whereas alkaline phosphatase activity declined.

#### 4. Discussion

In the previous study we have demonstrated the different developmental profile of  $11\beta$ HSD in ileum and colon and shown that corticosteroids stimulate  $11\beta$ HSD activity in immature and adult intestine [13]. This study is the first attempt to evaluate the possible role of  $11\beta$ HSD action in maturation of the intestine. Since

Table 1  
Kinetic parameters of ileal and colonic  $11\beta$ HSD of young rats

Tissue	$K_m$ (nM)	$V_{max}$ (pmol/min $\times$ mg protein)
Ileum	$59 \pm 10$	$66 \pm 6$
Distal colon	$81 \pm 14$	$8 \pm 1$

Kinetic study was performed in the presence of NAD as described in Section 2, 24-day-old rats were used. Values are means  $\pm$  S.E.M. of four separate experiments.




Fig. 3. Developmental changes of sucrase, Na,K-ATPase and alkaline phosphatase activities in jejunum, ileum and distal colon of intact rats during weaning. Each point is the mean  $\pm$  S.E.M. (sucrase:  $n = 10-17$ ; Na,K-ATPase:  $n = 6-8$ ; alkaline phosphatase:  $n = 8-10$ ).

Table 2

Effect of in vivo carbenoxolone administration on the activity of sucrase in jejunum and ileum

	Controls	Carbenoxolone
Jejunum	5.34 ± 0.79 [14]	6.71 ± 0.65 [15]
Ileum	1.61 ± 0.36 [15]	1.81 ± 0.28 [22]

The data are mean ± S.E.M., sucrase activity in  $\mu\text{mol glucose}/(\text{h} \times \text{mg protein})$ , numbers of animals in parentheses. Controls, 24-day-old rats that were i.p. treated with vehicle since day 12; carbenoxolone treated rats received the drug in a dose 60 mg/(kg body wt. day) since day 12 of life.

glucocorticoids are essential for maturation of intestine [1] and  $11\beta\text{HSD}$  regulates the availability of glucocorticoids in target tissues and cells [21–23],  $11\beta\text{HSD}$  might play an important role in the intestinal development. In addition,  $11\beta\text{HSD}$  has been proposed to protect mineralocorticoid receptors from binding glucocorticoids in mineralocorticoid-target tissue including distal colon [28] and immature distal colon has significantly increased  $\text{Na}^+$  absorption due to stimulation of electrogenic  $\text{Na}^+$  transport pathways by aldosterone [5]. However, the precise physiological role of  $11\beta\text{HSD}$  during development remains speculative.

We have demonstrated that the developmental profile of  $11\beta\text{HSD}$  activity in ileum is different from the profile in caecum and proximal or distal colon. The ileal developmental profile was comparable with that of  $11\beta\text{HSD1}$  isoform in liver [24], whereas the profile in the large intestinal segments was very similar to the profile in kidney [25] which is the major site of  $11\beta\text{HSD2}$  biosynthesis in adult animals [26]. The present results demonstrate clear-cut difference in cofactor preference in ileum and distal colon and indicate the existence of both isoforms in the intestine.  $K_m$  for rat liver  $11\beta\text{HSD1}$  is approximately  $1 \mu\text{M}$  [9] and NAD has no significant effect on its activity [27]. In contrast,  $11\beta\text{HSD2}$  in colonic T84 cells has  $K_m$  11 nM and clear preferences for NAD over NADP [19]. As we used substrate concentration 14.5 nM, our data seem to reflect the presence of  $11\beta\text{HSD2}$  in both ileum and distal colon. This conclusion is supported by the kinetic analysis (Table 1) which has demonstrated the presence of a high-affinity ileal and colonic  $11\beta\text{HSD2}$  similar to that recently reported in adult rat and human surface colonocytes [12]. The ‘mixed’

Table 3

Effect of in vivo carbenoxolone administration on the activities of alkaline phosphatase and Na,K-ATPase in ileum and distal colon of young rats

Enzyme	Treatment	Ileum	Distal colon
Alkaline phosphatase	Controls	17.1 ± 3.0 [9]	3.6 ± 0.6 [11]
	Carbenoxolone	17.7 ± 4.6 [11]	3.8 ± 0.5 [15]
Na,K-ATPase	Controls	6.55 ± 0.69 [14]	4.03 ± 0.59 [13]
	Carbenoxolone	6.22 ± 1.05 [15]	5.71 ± 0.42* [16]

Alkaline phosphatase in  $\mu\text{mol}$  of *p*-nitrophenol produced per hour and mg protein; Na,K-ATPase activity in  $\mu\text{mol}$  of phosphate produced per hour and mg protein. For further details see Table 2.

\*Significantly different from control animals ( $P < 0.05$ ).

cofactor preference probably reflects the presence of both isoforms of  $11\beta$ HSD in the intestine as was recently demonstrated in distal colon of adult rats [12]. According to these findings  $11\beta$ HSD2 is localized predominantly in the surface colonocytes which are responsible for  $\text{Na}^+$  transport and are target cells for mineralocorticoid action, whereas  $11\beta$ HSD1 is expressed predominantly in subepithelial cells. The question which is not yet resolved is the reversibility of colonic  $11\beta$ HSD1 isoform. Whorwood et al. [12] observed in isolated colonic subepithelial cells 11-oxo-reductase activity in the presence of 500 nM 11-dehydrocorticosterone but we have not seen any 11-oxo-reductase activity in slices of ileum and colon in the presence of 1.45  $\mu\text{M}$  11-dehydrocorticosterone [13,14]. In contrast, Marhefka et al. [20] detected 11-oxo-reductase activity in slices of duodenum, jejunum and ileum of adult rats. At present it is no explanation for these differences.

The results discussed above support the view that intestinal isoforms of  $11\beta$ HSD possess dehydrogenase activity and perhaps also oxo-reductase activity, i.e. they can increase or decrease glucocorticoid exposure to the glucocorticoid and mineralocorticoid receptors. However, the physiological significance of these processes remains to be fully defined. Postnatal maturation of the rodent intestine is characterized by significant modifications of enzyme status during the weaning period when it is modulated by corticosteroids [1,2]. To test the functional role of  $11\beta$ HSD in developing intestine, we have investigated the effect of  $11\beta$ HSD inhibitor carbenoxolone on the activity of three enzymes with large developmental changes during the weaning period which are sensitive to corticosteroids. These enzymes were sucrase, the activity of which substantially increases during weaning in jejunum and ileum [1]; alkaline phosphatase, whose activity decreases in ileum during weaning [16] and Na,K-ATPase whose activity rises postnatally in colon [14] and in ileum [15]. Even if sucrase and alkaline phosphatase activities are increased by glucocorticoids [1,29], we have not observed any significant changes of their activity in carbenoxolone-treated animals. In contrast, carbenoxolone administration significantly increased colonic but not ileal Na,K-ATPase. This is in agreement with the findings that, in immature rats, colonic Na,K-ATPase is increased by mineralocorticoids [31] and glucocorticoids [30] but not with the data of Zemelman et al. [15] who demonstrated the stimulatory effect of glucocorticoids on Na,K-ATPase in immature ileum. Similar effect of carbenoxolone on Na,K-ATPase was also observed in adult kidney [28,32]. The absence of the effect of carbenoxolone in ileum indicates that  $11\beta$ HSD has a protective effect on Na,K-ATPase only in mineralocorticoid-sensitive tissue. The activity of  $11\beta$ HSD in colonocytes of the immature rats may protect these cells from glucocorticoids and support the mineralocorticoid regulation of  $\text{Na}^+$  transport via  $\text{Na}^+$  channels and Na,K-ATPase [5,31]. The developmental increase of colonic Na,K-ATPase seems to be mineralocorticoid- and  $11\beta$ HSD-dependent whereas the glucocorticoid-dependent development of intestine does not seem to be influenced by this enzyme.

## Acknowledgements

This study was supported by Grants No. 305/94/1719 and No. 306/96/1289 from the Grant Agency of the Czech Republic. The authors thank Dr. J. Zicha (Institute of Physiology, Czech Acad. Sci., Prague) for reading the manuscript and valuable advice. The technical assistance of Mrs. R. Somolová is greatly appreciated.

## References

- [1] S.J. Henning, Functional development of the gastrointestinal tract, in: L.R. Johnson (Ed.), *Physiology of the Gastrointestinal Tract*, Raven Press, New York, 1987, pp. 285–300.
- [2] J. Pácha, Epithelial ion transport in the developing intestine, *Physiol. Res.* 42 (1993) 365–372.
- [3] S.J. Henning, Plasma concentrations of total and free corticosterone during development in the rat, *Amer. J. Physiol.* 235 (1978) E451–E456.
- [4] M. Kedinger, P.M. Simon, F. Raul, J.F. Grenier, K. Haffen, The effect of dexamethasone on the development of rat intestinal brush border enzymes in organ culture, *Dev. Biol.* 74 (1980) 9–11.
- [5] J. Pácha, I. Pohlová, P. Karen, Regulation of amiloride-sensitive  $\text{Na}^+$  transport in immature rat distal colon by aldosterone, *Pediatr. Res.* 38 (1995) 356–360.
- [6] C.R.W. Edwards, P.M. Stewart, The cortisol-cortisone shuttle and the apparent specificity of glucocorticoid and mineralocorticoid receptors, *J. Steroid Biochem. Mol. Biol.* 39 (1991) 859–865.
- [7] C. Monder, Corticosteroids, receptors, and the organ-specific functions of  $11\beta$ -hydroxysteroid dehydrogenase, *FASEB J.* 5 (1991) 3047–3054.
- [8] S.C. Low, K.E. Chapman, C. Edwards, J.R. Seckl, “Liver-type”  $11\beta$ -hydroxysteroid dehydrogenase cDNA encodes reductase but not dehydrogenase activity in intact mammalian COS-7 cells, *J. Mol. Endocrinol.* 13 (1994) 167–174.
- [9] V. Lakshmi, C. Monder, Purification and characterization of the corticosteroid  $11\beta$ -dehydrogenase component of the rat liver  $11\beta$ -hydroxysteroid dehydrogenase complex, *Endocrinology* 123 (1988) 2390–2398.
- [10] E. Rusvai, A. Naray-Fejes-Toth, A new isoform of  $11\beta$ -hydroxysteroid dehydrogenase in aldosterone target cells, *J. Biol. Chem.* 268 (1993) 10717–10720.
- [11] J. Pácha, I. Mikšík, Distribution of  $11\beta$ -hydroxysteroid dehydrogenase along the rat intestine, *Life Sci.* 54 (1994) 745–749.
- [12] C.B. Whorwood, M.L. Ricketts, P.M. Stewart, Epithelial cell localization of type2  $11\beta$ -hydroxysteroid dehydrogenase in rat and human colon, *Endocrinology* 135 (1994) 2533–2541.
- [13] J. Pácha, I. Mikšík,  $11\beta$ -Hydroxysteroid dehydrogenase in developing rat intestine, *J. Endocrinol.* 148 (1996) 561–566.
- [14] J. Pácha, J. Teisinger, M. Popp, K. Capek,  $\text{Na},\text{K}$ -ATPase and the development of  $\text{Na}^+$  transport in rat distal colon, *J. Membrane Biol.* 120 (1991) 201–210.
- [15] B.V. Zemelman, W.A. Walker, S. Chu, Expression and developmental regulation of  $\text{Na}^+, \text{K}^+$ -adenosine triphosphatase in the rat small intestine, *J. Clin. Invest.* 90 (1992) 1016–1022.
- [16] F. Moog, K.Y. Yeh, Intestinal alkaline phosphatase of the rat: Development and distribution of activity with phenylphosphate and  $\beta$ -glycerophosphate, *Comp. Biochem. Physiol.* 44B (1973) 657–666.
- [17] M.M. Bradford, A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal. Biochem.* 72 (1976) 248–254.
- [18] H. Murer, E. Amman, J. Biber, U. Hopfer, The surface membrane of the small intestinal epithelial cell. I, Localization of adenylyl cyclase, *Biochim. Biophys. Acta* 433 (1976) 509–519.
- [19] W.B. Reeves, NAD-dependent  $11\beta$ -hydroxysteroid dehydrogenase in cultured human colonic epithelial cells, *Am. J. Physiol.* 268 (1995) C1467–C1473.
- [20] A. Marhefka, U. Gross, K. Hierholzer, Segmental heterogeneity of corticosterone transformation along the intestinal tract. In vitro study with rat tissue, *Pflügers Arch.* 411 (1988) R106.

- [21] C.B. Whorwood, J.A. Franklyn, M.C. Sheppard, P.M. Stewart, Tissue localization of  $11\beta$ -hydroxysteroid dehydrogenase and its relationship to the glucocorticoid receptor, *J. Steroid Biochem. Mol. Biol.* 41 (1992) 21–28.
- [22] C.B. Whorwood, M.C. Sheppard, P.M. Stewart, Licorice inhibits  $11\beta$ -hydroxysteroid dehydrogenase messenger ribonucleic acid levels and potentiates glucocorticoid hormone action, *Endocrinology* 132 (1993) 2287–2292.
- [23] N. Page, N. Warriar, M.V. Govindan,  $11\beta$ -Hydroxysteroid dehydrogenase activity in human lung cells and transcription regulation by glucocorticoids, *Am. J. Physiol.* 267 (1994) L464–L474.
- [24] R. Ghraf, U. Vetter, J.M. Zandveld, H. Schiefers, Organ-specific ontogenesis of steroid hormone metabolizing enzyme activities in the rat, *Acta Endocrinol.* 79 (1975) 192–201.
- [25] M.P. Moisan, C.R.W. Edwards, J.R. Seckl, Ontogeny of  $11\beta$ -hydroxysteroid dehydrogenase in rat brain and kidney, *Endocrinology* 130 (1992) 400–404.
- [26] C. Monder, P.C. White,  $11\beta$ -hydroxysteroid dehydrogenase, *Vitamins and Hormones* 47 (1993) 187–271.
- [27] B.R. Walker, J.C. Campbell, B.C. Williams, C.R.W. Edwards, Tissue-specific distribution of the  $\text{NAD}^+$ -dependent isoform of  $11\beta$ -hydroxysteroid dehydrogenase, *Endocrinology* 131 (1992) 970–972.
- [28] C.B. Whorwood, M.L. Ricketts, P.M. Stewart, Regulation of sodium-potassium adenosine triphosphate subunit gene expression by corticosteroid and  $11\beta$ -hydroxysteroid dehydrogenase activity, *Endocrinology* 135 (1994) 901–910.
- [29] H. Pelichová, O. Koldovský, A. Heringová, V. Jirsová, J. Kraml, Postnatal changes of activity and electrophoretic pattern of jejunal and ileal nonspecific esterase and alkaline phosphatase of the rat, *Can. J. Biochem.* 45 (1967) 1375–1384.
- [30] P.J. Fuller, K. Verity, Colonic sodium-potassium adenosine triphosphatase subunit gene expression: Ontogeny and regulation by adrenocortical steroids, *Endocrinology* 127 (1990) 32–38.
- [31] Z. Zemanová, J. Pácha, Corticosteroid induction of renal and intestinal  $\text{K}^+$ -dependent *p*-nitrophenylphosphatase in young and adult rats, *Histochem. J.* 28 (1996) 625–634.
- [32] H. Tsuganezawa, M. Hayashi, Y. Fujii, Y. Yamaji, M. Iyori, T. Saruta, Corticosterone increases  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity in rat cortical collecting ducts with inhibition of  $11\beta$ -hydroxysteroid dehydrogenase, *Renal Physiol. Biochem.* 18 (1995) 66–72.